

B4

PCT

WORLD INTELLECTUAL PROPERTY ORGANIZATION
International Bureau

INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(51) International Patent Classification ⁷ : C12N 15/12, C07K 14/47, 16/18, G01N 33/566, C12Q 1/68, C12N 15/11, 15/62, A01K 67/027, A61K 38/00		A2	(11) International Publication Number: WO 00/58473
			(43) International Publication Date: 5 October 2000 (05.10.00)
(21) International Application Number: PCT/US00/08621		(72) Inventors; and	
(22) International Filing Date: 31 March 2000 (31.03.00)		(75) Inventors/Applicants (for US only): SHIMKETS, Richard, A. [US/US]; 191 Leete Street, West Haven, CT 06516 (US). LEACH, Martin [GB/US]; 884 School Street, Webster, MA 01570 (US).	
(30) Priority Data:		(74) Agent: ELRIFI, Ivor, R.; Mintz, Levin, Cohn, Ferris, Glovsky and Popeo, P.C., One Financial Center, Boston, MA 02111 (US).	
60/127,607	31 March 1999 (31.03.99)	US	
60/127,636	2 April 1999 (02.04.99)	US	
60/127,728	5 April 1999 (05.04.99)	US	
09/540,763	30 March 2000 (30.03.00)	US	
(63) Related by Continuation (CON) or Continuation-in-Part (CIP) to Earlier Applications		(81) Designated States: AE, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, CA, CH, CN, CR, CU, CZ, DE, DK, DM, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TR, TT, TZ, UA, UG, US, UZ, VN, YU, ZA, ZW, ARIPO patent (GH, GM, KE, LS, MW, SD, SL, SZ, TZ, UG, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE), OAPI patent (BF, BI, CF, CG, CI, CM, GA, GN, GW, ML, MR, NE, SN, TD, TG).	
US	60/127,607 (CIP)		
Filed on	31 March 1999 (31.03.99)		
US	60/127,636 (CIP)		
Filed on	2 April 1999 (02.04.99)		
US	60/127,728 (CIP)		
Filed on	5 April 1999 (05.04.99)		
US	09/540,763 (CIP)		
Filed on	30 March 2000 (30.03.00)		
(71) Applicant (for all designated States except US): CURAGEN CORPORATION [US/US]; 555 Long Wharf Drive, 11th Floor, New Haven, CT 06511 (US).		Published Without international search report and to be republished upon receipt of that report.	

(54) Title: NUCLEIC ACIDS INCLUDING OPEN READING FRAMES ENCODING POLYPEPTIDES; "ORFX"

(57) Abstract

The present invention provides open reading frames ORFX, encoding isolated polypeptides, as well as polynucleotides encoding ORFX and antibodies that immunospecifically bind to ORFX or any derivative, variant, mutant, or fragment of the ORFX polypeptides, polynucleotides or antibodies. The invention additionally provides methods in which the ORFX polypeptide, polynucleotide and antibody are used in detection and treatment of a broad range of pathological states, as well as to other uses.

FOR THE PURPOSES OF INFORMATION ONLY

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AL	Albania	ES	Spain	LS	Lesotho	SI	Slovenia
AM	Armenia	FI	Finland	LT	Lithuania	SK	Slovakia
AT	Austria	FR	France	LU	Luxembourg	SN	Senegal
AU	Australia	GA	Gabon	LV	Latvia	SZ	Swaziland
AZ	Azerbaijan	GB	United Kingdom	MC	Monaco	TD	Chad
BA	Bosnia and Herzegovina	GE	Georgia	MD	Republic of Moldova	TG	Togo
BB	Barbados	GH	Ghana	MG	Madagascar	TJ	Tajikistan
BE	Belgium	GN	Guinea	MK	The former Yugoslav Republic of Macedonia	TM	Turkmenistan
BF	Burkina Faso	GR	Greece			TR	Turkey
BG	Bulgaria	HU	Hungary	ML	Mali	TT	Trinidad and Tobago
BJ	Benin	IE	Ireland	MN	Mongolia	UA	Ukraine
BR	Brazil	IL	Israel	MR	Mauritania	UG	Uganda
BY	Belarus	IS	Iceland	MW	Malawi	US	United States of America
CA	Canada	IT	Italy	MX	Mexico	UZ	Uzbekistan
CF	Central African Republic	JP	Japan	NE	Niger	VN	Viet Nam
CG	Congo	KE	Kenya	NL	Netherlands	YU	Yugoslavia
CH	Switzerland	KG	Kyrgyzstan	NO	Norway	ZW	Zimbabwe
CI	Côte d'Ivoire	KP	Democratic People's Republic of Korea	NZ	New Zealand		
CM	Cameroon	KR	Republic of Korea	PL	Poland		
CN	China			PT	Portugal		
CU	Cuba	KZ	Kazakhstan	RO	Romania		
CZ	Czech Republic	LC	Saint Lucia	RU	Russian Federation		
DE	Germany	LI	Liechtenstein	SD	Sudan		
DK	Denmark	LK	Sri Lanka	SE	Sweden		
EE	Estonia	LR	Liberia	SG	Singapore		

What is claimed is:

1. An isolated nucleic acid molecule encoding a polypeptide comprising an amino acid sequence that is at least 85% identical to a polypeptide including an amino acid sequence selected from the group consisting of SEQ ID NO:2*n*, wherein *n* is any integer 1-3161, or the complement thereof.
2. The isolated nucleic acid molecule of claim 1, said molecule hybridizing under stringent conditions to a nucleic acid sequence complementary to a nucleic acid molecule comprising the sequence of nucleotides selected from the group consisting of SEQ ID NO:2*n*, wherein *n* is any integer 1-3161, or the complement thereof.
3. The isolated nucleic acid molecule of claim 1, said molecule encoding a polypeptide comprising the amino acid sequence selected from the group consisting of SEQ ID NO: 2*n*, wherein *n* is any integer 1-3161, or an amino acid sequence comprising one or more conservative substitutions in the amino acid sequence selected from the group consisting of SEQ ID NO: 2*n*.
4. The isolated nucleic acid molecule of claim 1, wherein said molecule encodes a polypeptide comprising the amino acid sequence selected from the group consisting of SEQ ID NO: 2*n*, wherein *n* is any integer 1-3161.
5. The isolated nucleic acid molecule of claim 1, wherein said molecule comprises the sequence of nucleotides selected from the group consisting of SEQ ID NO:2*n*-1, wherein *n* is any integer 1-3161, or the complement thereof.
6. An oligonucleotide less than 100 nucleotides in length and comprising at least one contiguous nucleotide selected from the group consisting of SEQ ID NO:2*n*-1, wherein *n* is any integer 1-3161, or the complement thereof.
7. A vector comprising the nucleic acid molecule of claim 1.

8. The vector of claim 7, wherein said vector is an expression vector.
9. A host cell comprising the isolated nucleic acid molecule of claim 1.
10. A substantially purified polypeptide comprising an amino acid sequence at least 80% identical to a polypeptide comprising the amino acid sequence selected from the group consisting of SEQ ID NO: 2*n*, wherein *n* is any integer 1-3161.
11. The polypeptide of claim 10, wherein said polypeptide comprises the amino acid sequence selected from the group consisting of SEQ ID NO: 2*n*, wherein *n* is any integer 1-3161.
12. An antibody that selectively binds to the polypeptide of claim 10.
13. A pharmaceutical composition comprising a therapeutically or prophylactically effective amount of a therapeutic selected from the group consisting of:
 - a) the nucleic acid of claim 1;
 - b) the polypeptide of claim 10; and
 - c) the antibody of claim 12;and a pharmaceutically acceptable carrier.
14. A kit comprising in one or more containers, a therapeutically or prophylactically effective amount of the pharmaceutical composition of claim 13.
15. A method of producing the polypeptide of claim 10, said method comprising culturing the host cell of claim 9 under conditions in which the nucleic acid molecule is expressed.
16. A method of detecting the presence of the polypeptide of claim 10 in a sample, comprising contacting the sample with a compound that selectively binds to said polypeptide under conditions allowing the formation of a complex between said polypeptide and said

compound, and detecting said complex, if present, thereby identifying said polypeptide in said sample.

17. A method of detecting the presence of a nucleic acid molecule of claim 1 in a sample, the method comprising contacting the sample with a nucleic acid probe or primer that selectively binds to the nucleic acid molecule and determining whether the nucleic acid probe or primer bound to the nucleic acid molecule of claim 1 is present in the sample.

18. A method for modulating the activity of the polypeptide of claim 10, the method comprising contacting a cell sample comprising the polypeptide of claim 10 with a compound that binds to said polypeptide in an amount sufficient to modulate the activity of the polypeptide.

19. The use of a therapeutic in the manufacture of a medicament for treating a syndrome associated with a ORFX-associated disorder, wherein said therapeutic is selected from the group consisting of:

- a) the nucleic acid of claim 1;
- b) the polypeptide of claim 10; and
- c) the antibody of claim 12.

20. A method for screening for a modulator of activity or of latency or predisposition to an ORFX-associated disorder, said method comprising:

- a) contacting a test compound with the polypeptide of claim 10; and
- b) determining if said test compound binds to said polypeptide,

wherein binding of said test compound to said polypeptide indicates the test compound is a modulator of activity or of latency or predisposition to an ORFX-associated disorder.

21. A method for screening for a modulator of activity or of latency or predisposition to an ORFX-associated disorder, said method comprising:

- a) administering a test compound to a test subject at an increased risk ORFX-associated disorder, wherein said test subject recombinantly expresses a polypeptide encoded by the nucleotide of claim 1;

- b) measuring expression the activity of said protein in said test subject;
- c) measuring the activity of said protein in a control subject that recombinantly expresses said protein and is not at increased risk for an ORFX-associated disorder; and
- d) comparing expression of said protein in said test subject and said control subject, wherein a change in the activity of said protein in said test subject relative to said control subject indicates the test compound is a modulator or of latency of predisposition to an ORFX-associated disorder.

22. The method of claim 20, wherein said test animal is a recombinant test animal that expresses a test protein transgene or expresses said transgene under the control of a promoter at an increased level relative to a wild-type test animal, and wherein said promoter is not the native gene promoter of said transgene.

23. A method for determining the presence of or predisposition to a disease associated with altered levels of a polypeptide of claim 11 in a subject, the method comprising:

- a) measuring the amount of the polypeptide in a sample from said subject; and
- b) comparing the amount of said polypeptide in step (a) to the amount of the polypeptide present in a control sample,

wherein an alteration in the level of the polypeptide in step (a) as compared to the control sample indicates the presence of or predisposition to a disease in said subject.

24. The method of claim 23, wherein said subject is a human.

25. A method for determining the presence of or predisposition to a disease associated with altered levels the nucleic acid molecule of claim 1 in a subject, the method comprising:

- a) measuring the amount of the nucleic acid in a sample from the mammalian subject; and
- b) comparing the amount of said nucleic acid in step (a) to the amount of the nucleic acid present in a control sample,

wherein an alteration in the level of the nucleic acid in step (a) as compared to the control sample indicates the presence of or predisposition to said disease in said subject.

26. The method of claim 25, wherein said subject is a human.

27. A method of treating or preventing a pathological condition associated with an ORFX-associated disorder in a subject, the method comprising administering to said subject a polypeptide of claim 10 in an amount sufficient to alleviate or prevent said pathological condition.


28. The method of claim 27, wherein said subject is a human.

29. A method of treating or preventing a pathological condition associated with an ORFX-associated disorder in a subject, the method comprising administering to said subject a nucleic acid molecule of claim 1 in an amount sufficient to alleviate or prevent said pathological condition.

30. The method of claim 29, wherein said subject is a human.

31. A method of treating or preventing a pathological condition associated with an ORFX-associated disorder in a subject, the method comprising administering to said subject an antibody of claim 12 in an amount sufficient to alleviate or prevent said pathological condition.

32. The method of claim 31, wherein said subject is a human.



<213> Homo sapiens

<400> 360

```

Thr Arg Thr Asp Lys Arg Pro Val Met Ala Asp Leu Arg Glu Ser Gly
 1           5           10           15
Ala Ile Glu Gln Asp Ala Asp Met Ile Val Phe Ile Tyr Arg Asp Asp
          20           25           30
Tyr Tyr Asn Lys Glu Asn Ser Pro Asp Lys Gly Leu Ala Glu Ile Ile
      35           40           45
Ile Gly Lys His Arg Gly Gly Pro Thr Gly Ser Cys Lys Leu Lys Phe
      50           55           60
Phe Gly Glu Tyr Thr Arg Phe Asp Asn Leu Ala His Asn Ser Val Gly
65           70           75           80
Ser Phe Glu

```

<210> 361

<211> 453

<212> DNA

<213> Homo sapiens

<400> 361

```

gctttgcagg aggaatatctc tatctctggc tgcaagatga ggctgagcta cctgagcagc
60
cggaccccctg gctacaaatc tgtcctgagg atcagcctca ccaccccgac catccccttc
120
aacctcatga aggtgcacct catggtagcg gtggaggccc gccctcttcag gaagtgttcc
180
gtgagcagccc cagacctgtc ctattatttc atttgggaca agacagacgt ctacaaccag
240
aagggtgtttg ggctttcaga agcctttgtt tccgtgggtt atgaatatga atcctgccca
300
gatctaattcc tgtgggaaaa aagaacaaca gtgctgcagg gctatgaaat tgacgcgtcc
360
aagcttggag gatggagcct agacaaacat catgcctca acattcaag tggcatcctg
420
cacaaagga atggngagaa ccagttttgtg tct
453

```

<210> 362

<211> 151

<212> PRT


<213> Homo sapiens

<400> 362

```

Ala Leu Gln Glu Glu Ile Ser Ile Ser Gly Cys Lys Met Arg Leu Ser
 1           5           10           15
Tyr Leu Ser Ser Arg Thr Pro Gly Tyr Lys Ser Val Leu Arg Ile Ser
      20           25           30
Leu Thr His Pro Thr Ile Pro Phe Asn Leu Met Lys Val His Leu Met
      35           40           45
Val Ala Val Glu Gly Arg Leu Phe Arg Lys Trp Phe Ala Ala Ala Pro
      50           55           60
Asp Leu Ser Tyr Tyr Phe Ile Trp Asp Lys Thr Asp Val Tyr Asn Gln

```



```

65              70              75              80
Lys Val Phe Gly Leu Ser Glu Ala Phe Val Ser Val Gly Tyr Glu Tyr
85
Glu Ser Cys Pro Asp Leu Ile Leu Trp Glu Lys Arg Thr Val Leu
100
Gln Gly Tyr Glu Ile Asp Ala Ser Lys Leu Gly Gly Trp Ser Leu Asp
115
Lys His His Ala Leu Asn Ile Gln Ser Gly Ile Leu His Lys Gly Asn
130
Gly Glu Asn Gln Phe Val Ser
145
150

```

<210> 363
 <211> 502
 <212> DNA
 <213> Homo sapiens

```

<400> 363
ggtaacacaaa aagtttgcca cagtattcac actccaggctc tccataaacc ttccagatcc
60
gctcacacaaa gctgggtgttc atttgettct tctgtaaaact gtccaggacc ttcattgaaag
120
cggtgatgcc tgaccggtgc tcaggggagc ctttgcaaga gtcaggctga tgtgtgatgg
180
tgtccccacc accagctact ggagggagga ggtctgaggg ctcagctggg ttgacctga
240
gacacctgct gggatctggg tcaccagctg aaagcacagc catgttctgc ccttccccta
300
gggggctctg ggcgccatgg ctttctctgat ctgaccacgc actctgggccc ttggacagca
360
gtagtgtgat cacttcacct tgcgtctgga ctgagcttct gtgctgcatg tctgggggct
420
tctcaggagc agcatgagcc tctcgggagg aggtatcatt ttcaacaaa aaatcatctg
480
aaaccacctc ttgagaatgc ag
502


```

<210> 364
 <211> 136
 <212> PRT
 <213> Homo sapiens

```

<400> 364
Met Gln His Arg Ser Ser Val Gln Thr Gln Gly Glu Val Ile Thr Leu
1              5              10              15
Leu Leu Ser Lys Ala Gln Ser Ala Gly Ser Asp Gln Glu Ser His Gly
20
Ala Gln Ser Pro Leu Gly Glu Gly Gln Asn Met Ala Val Leu Ser Ala
35
Gly Asp Pro Asp Pro Ser Arg Cys Leu Arg Ser Asn Pro Ala Glu Ala
50
Ser Asp Leu Leu Pro Pro Val Ala Gly Gly Asp Thr Ile Thr His
65
Gln Pro Asp Ser Cys Lys Ala Ala Pro Glu His Arg Ser Gly Ile Thr
70              75              80

```



ID HSM010618 standard; RNA; EST; 718 BP.

XX
AC AL045768;

XP-002237532

XX
SV AL045768.1

XX
DT 12-MAR-1999 (Rel. 59, Created)
DT 20-FEB-2000 (Rel. 62, Last updated, Version 4)

XX
DE Homo sapiens mRNA; EST DKFZp434F206_r1 (from clone DKFZp434F206)

XX
KW EST; expressed sequence tag.

XX
OS Homo sapiens (human)

OC Eukaryota; Metazoa; Chordata; Craniata; Vertebrata; Euteleostomi; Mammalia;
OC Eutheria; Primates; Catarrhini; Hominidae; Homo.

XX
RN [1]

RP 1-718

RA Koehler K., Beyer A., Mewes H.W., Gassenhuber J., Wiemann S.;

RT ;

RL Submitted (18-FEB-2000) to the EMBL/GenBank/DBJ databases.

RL MIPS, Am Klopferspitz 18a, D-82152 Martinsried, GERMANY

XX
DR RZPD; DKFZp434F206; DKFZp434F206.

DR UNILIB; 1752; 1752.

XX
CC This is the 5' sequence of the clone insert

CC Clone from S. Wiemann, Molecular Genome Analysis, German Cancer
CC Research Center (DKFZ); Email s.wiemann@dkfz-heidelberg.de;
CC sequenced by BMFZ (Biomedical Research Center at the Charite,
CC Berlin/Germany) within the cDNA sequencing consortium of the
CC German Genome Project.

CC s1 sequence also available.

CC This clone (DKFZp434F206) is available at the RZPD in Berlin.

CC Please contact the RZPD: Ressourcenzentrum, Heubnerweg 6,

CC 14059 Berlin-Charlottenburg, GERMANY; Email: clone@rzd.de

XX
FH Key Location/Qualifiers

FH
FT source

1..718

/db_xref="taxon:9606"

/db_xref="RZPD:DKFZp434F206"

/db_xref="UNILIB:1752"

/organism="Homo sapiens"

/clone="DKFZp434F206"

/clone_lib="434 (synonym: htes3). Vector pSport1; host

DH10B; sites NotI + SalI"

/dev_stage="adult"

/tissue_type="testis"

XX
SQ

Sequence 718 BP; 183 A; 196 C; 198 G; 141 T; 0 other;
gcttcacagc agatgtttaac agctggctgc tcaacctttgg attccagcta cacacagtga 60
tccctgtgta tcccaaacca gacatggatg ccatggaaac ctctacagag cttatccaca 120
cacagatgaa aacgcaggag tgggacaaca gcaagtctat cctcggggta caggtggaag 180
tacagaagca gctcaaggcc ttgttcacct tagaacggtt tgaccagctc tatggctcca 240
caatcaccaag ctgccagcag gctccaaaga ccaagaagtt tgcatccagc ggctcagttc 300
ttggcgaagg ggtcaagttt gcttgaagg atggccgagt gaccacagac atcatcagtg 360
tgcccaatga ggaatggcga aggggttgcg ccatcttgaa ccatgccac tacctagaga 420
acctgacctt caccattgat ggggtggata cccattact ttgtgaaacca ggaccttcag 480
aaggtgacct ggccatcctg ggccctcagtg gggggcggcg aacctctgag aaatgggttca 540
acgtcacctg gtcccaagtc aacacagatc ttaatggcag gactagacgc tacacagaca 600
tccagctcca gtacggggca ctgtgcttga acacacgcta cgggacaacg ttggaatgag 660
agaaggcagc ggtcctcagg ctgtcccgcc agagagccgt gcgccaagcg tgggcccc 718

